VASCULITIS DAMAGE INDEX (VDI)

This is for recording organ damage that has occurred in patients <u>since the onset of vasculitis</u> Patients often have co-morbidity before they develop vasculitis, **which must not be scored** Record features of active disease using the Birmingham Vasculitis Activity Score (BVAS)

A new patient should *usually have a VDI score of zero*, unless:

(a) they have had vasculitis for more than three months of onset of disease. and

(b) the damage has developed or become worse since the onset of vasculitis

1. MusculoskeletalNoNone□Significant muscle atrophy or weakness□Deforming/erosive arthritis□Osteoporosis/vertebral collapse□Avascular necrosis□Osteomyelitis□2. Skin/Mucous membranes□	Yes 0 0 0 0 0	Name Trial Number Date Centre 7. Peripheral vascular disease None	No	Yes
Significant muscle atrophy or weakness Deforming/erosive arthritis Osteoporosis/vertebral collapse Avascular necrosis Osteomyelitis 2. Skin/Mucous membranes	0 0 0	Date Centre 7. Peripheral vascular disease None		Yes
Deforming/erosive arthritis Osteoporosis/vertebral collapse Avascular necrosis Osteomyelitis 2. Skin/Mucous membranes	0 0 0	Centre 7. Peripheral vascular disease None		Yes
Osteoporosis/vertebral collapse Avascular necrosis Osteomyelitis 2. Skin/Mucous membranes	0 0	None		Yes
Avascular necrosis Osteomyelitis 2. Skin/Mucous membranes	0	None		res
Osteomyelitis 2. Skin/Mucous membranes				
2. Skin/Mucous membranes	0			\circ
		Absent pulses in one limb		0 0
		2 nd episode of absent pulses in one limb		
Alemania	0	Major vessel stenosis		0
Alopecia	0	Claudication >3 mths		0
Cutaneous ulcers	0	Minor tissue loss		0
Mouth ulcers	0	Major tissue loss		0
3. Ocular		Subsequent major tissue loss		0
None 🗆	~	Complicated venous thrombosis		0
Cataract	0	8. Gastrointestinal	_	
Retinal change	0	None		~
Optic atrophy	0	Gut infarction/resection		0
Visual impairment/diplopia	0	Mesenteric insufficiency/pancreatitis		0
Blindness in one eye	0	Chronic peritonitis		0
Blindness in second eye	0	Oesophageal stricture/surgery		0
Orbital wall destruction	0	9. Renal	_	
4. ENT		None		~
None 🛛	-	Estimated/measured GFR ≤ 50%		0
Hearing loss	0	Proteinuria ≥ 0.5g/24hr		0
Nasal blockage/chronic discharge/crusting	0	End stage renal disease		0
Nasal bridge collapse/septal perforation	0	10. Neuropsychiatric	_	
Chronic sinusitis/radiological damage	0	None		-
Subglottic stenosis (no surgery)	0	Cognitive impairment		0
Subglottic stenosis (with surgey)	0	Major psychosis		0
5. Pulmonary		Seizures		0
None		Cerebrovascular accident		0
Pulmonary hypertention	0	2 nd cerebrovascular accident		0
Pulmonary fibrosis	0	Cranial nerve lesion		0
Pulmonary infarction	0	Peripheral neuropathy		0
Pleural fibrosis	0	Transverse myelitis		0
Chronic asthma	0	11. Other		
Chronic breathlessness	0	None		
Impaired lung function	0	Gonadal failure		0
6. Cardiovascular		Marrow failure		0
None 🛛		Diabetes		0
Angina angioplasty	0	Chemical cystitis		0
Myocardial infarction	0	Malignancy		0
Subsequent myocardial infarction	0	Other		0
Cardiomyopathy	0	Total VDI Score. Record the number of po	sitive	
Valvular disease	0	Items (1 point for each). The VDI score can		
Pericaritis ≥ 3 mths or pericardectomy	0	either increase or remain the same over time.		
Diastolic BP ≥ 95 or requiring	0	Remember to carry forward any previous items of damage.		
antihypertensives			L	

VDI Modified from Exley AR, Bacon PA, Luqmani et al (1997) Development and initial validation of the VDI ... Arthritis Rheum 40: 371-380